

# Vietnam Journal of MATHEMATICS

Formerly *Tạp chí Toán học (Journal of Mathematics)*

---

**Volume 33, Number 4 December 2005**

---

**Editor-in-Chief**

Nguyen Khoa Son  
Institute of Mathematics, Hanoi

**Deputy Editors-in-Chief**

Pham Ky Anh,  
Vietnam National University, Hanoi

Nguyen Tu Cuong  
Institute of Mathematics, Hanoi

Hoang Xuan Phu  
Institute of Mathematics, Hanoi

**Editorial Board**

- Hans George Bock, Universität Heidelberg  
Markus Brodmann, Universität Zürich  
Duong Minh Duc, University of Ho Chi Minh City  
Dinh Dung, Information Technology Institute, VNU, Hanoi  
Karl-Heinz Hoffmann, Caesar, Bonn  
Nguyễn Hữu Việt Hưng, Vietnam National University, Hanoi  
Masami Ito, Kyoto Sangyo University, Kyoto  
Ha Huy Khoai, Institute of Mathematics, Hanoi  
Lê Hai Khôi, Institute of Information Technology, Hanoi  
Pham The Long, Le Qui Don University, Hanoi  
Charles A. Micchelli, State University of New York, New York  
Tran Van Nhung, Ministry of Education and Training, Hanoi  
Shum Kar Ping, The Chinese University of Hong Kong  
J. P. Ramis, Université Paul Sabatier, Toulouse  
Vladimir N. Temlyakov, University of South Carolina, Columbia  
Do Duc Thai, Vietnam National University, Hanoi  
Dang Hung Thang, Vietnam National University, Hanoi  
Nguyen Quoc Thang, Institute of Mathematics, Hanoi  
Dao Trong Thi, Vietnam National University, Hanoi  
Do Long Van, Institute of Mathematics, Hanoi  
Tran Duc Van, Institute of Mathematics, Hanoi  
Nguyen Thanh Van, Université Paul Sabatier, Toulouse  
Ha Huy Vui, Institute of Mathematics, Hanoi  
Robert Wisbauer, Heinrich - Heine - Universität, Düsseldorf  
Steven Zucker, Johns Hopkins University, Baltimore

VIETNAMESE ACADEMY OF  
SCIENCE AND TECHNOLOGY  
&  
VIETNAM MATHEMATICAL SOCIETY



Springer

## Central Limit Theorem for Functional of Jump Markov Processes

Nguyen Van Huu, Vuong Quan Hoang, and Tran Minh Ngoc

*Department of Mathematics  
Hanoi National University, 334 Nguyen Trai Str., Hanoi, Vietnam*

Received February 8, 2005

Revised May 19, 2005

**Abstract.** In this paper some conditions are given to ensure that for a jump homogeneous Markov process  $\{X(t), t \geq 0\}$  the law of the integral functional of the process:  $T^{-1/2} \int_0^T \varphi(X(t))dt$ , converges to the normal law  $N(0, \sigma^2)$  as  $T \rightarrow \infty$ , where  $\varphi$  is a mapping from the state space  $E$  into  $\mathbb{R}$ .

### 1. Introduction

The central limit theorem is a subject investigated intensively by many well-known probabilists such as Linderberg, Chung,.... The results concerning central limit theorems, the iterated logarithm law, the lower and upper bounds of the moderate deviations are well understood for independent random variable sequences and for martingales but less is known for dependent random variables such as Markov chains and Markov processes.

The first result on central limit for functionals of stationary Markov chain with a finite state space can be found in the book of Chung [5]. A technical method for establishing the central limit is the regeneration method. The main idea of this method is to analyse the Markov process with arbitrary state space by dividing it into independent and identically distributed random blocks between visits to fixed state (or atom). This technique has been developed by Athreya - Ney [2], Nummelin [10], Meyn - Tweedie [9] and recently by Chen [4].

The technical method used in this paper is based on central limit for martingales and ergodic theorem. The paper is organized as follows:

In Sec. 2, we shall prove that for a positive recurrent Markov sequence

$\{X_n, n \geq 0\}$  with Borel state space  $(E, \mathcal{B})$  and for  $\varphi : E \rightarrow \mathbb{R}$  such that

$$\varphi(x) = f(x) - Pf(x) = f(x) - \int_E f(y)P(x, dy)$$

with  $f : E \rightarrow \mathbb{R}$  such that  $\int_E f^2(x)\Pi(dx) < \infty$ , where  $P(x, \cdot)$  is the transition probability and  $\Pi(\cdot)$  is the stationary distribution of the process, the distribution of  $n^{-1/2} \sum_{i=1}^n \varphi(X_i)$  converges to the normal law  $N(0, \sigma^2)$  with  $\sigma^2 = \int_E (\varphi^2(x) + 2\varphi(x)Pf(x))\Pi(dx)$ .

The central limit theorem for the integral functional  $T^{-1/2} \int_0^T \varphi(X(t))dt$  of jump Markov process  $\{X(t), t \geq 0\}$  will be established and proved in Sec. 3.

Some examples will be given in Sec. 4.

It is necessary to emphasize that the conditions for normal asymptoticity of  $n^{-1/2} \sum_{i=1}^n \varphi(X_i)$  is the same as in [8] but they are not equivalent to the ones established in [10, 11]. The results on the central limit for jump Markov processes obtained in this paper are quite new.

## 2. Central Limit for the Functional of Markov Sequence

Let us consider a Markov sequence  $\{X_n, n \geq 0\}$  defined on a basic probability space  $(\Omega, \mathcal{F}, P)$  with the Borel state space  $(E, \mathcal{B})$ , where  $\mathcal{B}$  is the  $\sigma$ -algebra generated by the countable family of subsets of  $E$ . Suppose that  $\{X_n, n \geq 0\}$  is homogeneous with transition probability

$$P(x, A) = P(X_{n+1} \in A | X_n = x), \quad A \in \mathcal{B}.$$

We have the following definitions

**Definition 2.1.** Markov process  $\{X_n, n \geq 0\}$  is said to be irreducible if there exists a  $\sigma$ -finite measure  $\mu$  on  $(E, \mathcal{B})$  such that for all  $A \in \mathcal{B}$

$$\mu(A) > 0 \text{ implies } \sum_{n=1}^{\infty} P^n(x, A) > 0, \quad \forall x \in E$$

where

$$P^n(x, A) = P(X_{m+n} \in A | X_m = x).$$

The measure  $\mu$  is called irreducible measure.

By Proposition 2.4 of Nummelin [10], there exists a maximum irreducible measure  $\mu^*$  possessing the property that if  $\mu$  is any irreducible measure then  $\mu \ll \mu^*$ .

**Definition 2.2.** Markov process  $\{X_n, n \geq 0\}$  is said to be recurrent if

$$\sum_{n=1}^{\infty} P^n(x, A) = \infty, \quad \forall x \in E, \forall A \in \mathcal{B} : \mu^*(A) > 0.$$

The process is said to be Harris recurrent if

$$P_x(X_n \in A \text{ i.o.}) = 1.$$

Let us notice that a process which is Harris recurrent is also recurrent.

**Theorem 2.1.** *If  $\{X_n, n \geq 0\}$  is recurrent then there exists a uniquely invariant measure  $\Pi(\cdot)$  on  $(E, \mathcal{B})$  (up to constant multiples) in the sense*

$$\Pi(A) = \int_E \Pi(dx)P(x, A), \quad \forall A \in \mathcal{B}, \tag{1}$$

or equivalently

$$\Pi(\cdot) = \Pi P(\cdot). \tag{2}$$

(see Theorem 10.4.4 of Meyn-Tweedie, [9]).

**Definition 2.3.** *A Markov sequence  $\{X_n, n \geq 0\}$  is said to be positive recurrent (null recurrent) if the invariant measure  $\Pi$  is finite (infinite).*

For a positive recurrent Markov sequence  $\{X_n, n \geq 0\}$ , its unique invariant probability measure is called stationary distribution and is denoted by  $\Pi$ . Hereafter we always denote the stationary distribution of Markov sequence  $\{X_n, n \geq 0\}$  by  $\Pi$  and if  $\nu$  is the initial distribution of Markov sequence then  $P_\nu(\cdot), E_\nu(\cdot)$  are denoted for probability and expectation operator responding to  $\nu$ . In particular,  $P_\nu(\cdot), E_\nu(\cdot)$  are replaced by  $P_x(\cdot), E_x(\cdot)$  if  $\nu$  is the Dirac measure at  $x$ .

We have the following ergodic theorem:

**Theorem 2.2.** *If Markov sequence  $\{X_n, n \geq 0\}$  possesses the unique invariant distribution  $\Pi$  such that*

$$P(x, \cdot) \ll \Pi(\cdot), \quad \forall x \in E, \tag{3}$$

then  $\{X_n, n \geq 0\}$  is metrically transitive when initial distribution is the stationary distribution. Further, for any measurable mapping  $\varphi : E \times E \rightarrow \mathbb{R}$  such that  $E_\Pi|\varphi(X_0, X_1)| < \infty$ , with probability one

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{k=0}^{n-1} \varphi(X_k, X_{k+1}) = E_\Pi \varphi(X_0, X_1) \tag{4}$$

and the limit does not depend on the initial distribution. (See Theorem 1.1 from Patrick Billingsley [3]).

The following notations will be used in this paper: For a measurable mapping  $\varphi : E \rightarrow \mathbb{R}$  we denote

$$\begin{aligned} \Pi\varphi &= \int_E \varphi(x)\Pi(dx), \quad P\varphi(x) = \int_E \varphi(y)P(x, dy) = E(\varphi(X_{n+1})|X_n = x), \\ P^n\varphi(x) &= \int_E \varphi(y)P^n(x, dy) = E(\varphi(X_{n+m})|X_m = x). \end{aligned}$$

For the countable state space  $E = \{1, 2, \dots\}$  we denote

$$\begin{aligned} P_{ij} &= P(i, \{j\}) = P(X_{n+1} = j | X_n = i), P_{ij}^{(n)} \\ &= P^n(i, \{j\}) = P(X_{m+n} = j | X_n = i) \\ \pi_j &= \Pi(\{j\}), P = [P_{ij}, i, j \in E], P^{(n)} = [P_{ij}^{(n)}, i, j \in E] = P^n. \end{aligned}$$

Then

$$\Pi\varphi = \sum_{j \in E} \varphi(j)\pi_j, P\varphi(j) = \sum_{k \in E} \varphi(k)P_{jk}, P^n\varphi(j) = \sum_{k \in E} \varphi(k)P_{jk}^{(n)}.$$

If the distribution of random variable  $Y_n$  converges to the normal distribution  $N(\mu, \sigma^2)$  then we denote  $\xrightarrow{\mathcal{L}} N(\mu, \sigma^2)$ . The indicator function of a set  $A$  is denoted by  $\mathbf{1}_A$ , where

$$\mathbf{1}_A(\omega) = \begin{cases} 1, & \text{if } \omega \in A \\ 0, & \text{else.} \end{cases}$$

Finally, the mapping  $\varphi : E = \{1, 2, \dots\} \rightarrow \mathbb{R}$  is denoted by column vector  $\varphi = (\varphi(1), \varphi(2), \dots)^T$ .

The main result of this section is to establish the conditions for

$$n^{-1/2} \sum_{k=1}^n \varphi(X_k) \xrightarrow{\mathcal{L}} N(\mu, \sigma^2).$$

We need a central limit theorem for martingale differences as follows

**Theorem 2.3.** (Central limit theorem for martingale differences) *Suppose that  $\{u_k, k \geq 0\}$  is a sequence of martingale differences defined on a probability space  $(\Omega, \mathcal{F}, P)$  corresponding to a filter  $\{\mathcal{F}_k, k \geq 0\}$ , i.e.,  $E(u_{k+1} | \mathcal{F}_k) = 0, k = 0, 1, 2, \dots$ . Further, assume that the following conditions are satisfied*

$$(A_1) \quad n^{-1} \sum_{k=1}^n E(u_k^2 | \mathcal{F}_{k-1}) \xrightarrow{P} \sigma^2,$$

$$(A_2) \quad n^{-1} \sum_{k=1}^n E(u_k^2 \mathbf{1}_{\{|u_k| \geq \varepsilon \sqrt{n}\}} | \mathcal{F}_{k-1}) \xrightarrow{P} 0, \text{ for each } \varepsilon > 0 \text{ (the conditional Lindeberg's condition).}$$

Then

$$n^{-1/2} \sum_{k=1}^n u_k \xrightarrow{\mathcal{L}} N(0, \sigma^2). \tag{5}$$

(see Corollary of Theorem 3.2, [7]).

*Remark 1.* Theorem 2.3 remains valid for  $\{u_k, k \geq 0\}$  being a  $m$ -dimensional martingale differences where the condition  $(A_1)$  is replaced by

$$n^{-1} \sum_{k=1}^n \text{Var} (u_k | \mathcal{F}_{k-1}) \xrightarrow{P} \sigma^2 = [\sigma_{ij}, i, j = 1, 2, \dots, m]$$

with

$$\text{Var}(u_k | \mathcal{F}_{k-1}) = [E(u_{ik}u_{jk} | \mathcal{F}_{k-1}), i, j = 1, 2, \dots, m].$$

We shall prove the following theorem.

**Theorem 2.4.** (Central limit theorem for functional of Markov sequence) *Suppose that the following conditions hold:*

(H<sub>1</sub>) *The Markov sequence {X<sub>n</sub>, n ≥ 0} is positive recurrent with the transition probability P(x, .) and the unique stationary distribution Π(.) satisfying the condition (3).*

(H<sub>2</sub>) *The mapping φ : E → ℝ can be represented in the form*

$$\varphi(x) = f(x) - Pf(x), \quad x \in E, \tag{6}$$

where f : E → ℝ is measurable and Πf<sup>2</sup> < ∞.

Then

$$n^{-1/2} \sum_{k=1}^n \varphi(X_k) \xrightarrow{\mathcal{L}} N(0, \sigma^2) \tag{7}$$

for any initial distribution, where

$$\sigma^2 = \Pi(f^2 - (Pf)^2) = \Pi(\varphi^2 + 2\varphi Pf). \tag{8}$$

*Proof.* We have

$$\begin{aligned} n^{-1/2} \sum_{k=1}^n \varphi(X_k) &= n^{-1/2} \sum_{k=1}^n [f(X_k) - Pf(X_k)] \\ &= n^{-1/2} \sum_{k=1}^n [f(X_k) - Pf(X_{k-1})] + n^{-1/2} \sum_{k=1}^n Pf(X_{k-1}) - n^{-1/2} \sum_{k=1}^n Pf(X_k) \\ &= n^{-1/2} \sum_{k=1}^n u_k + n^{-1/2} [Pf(X_0) - Pf(X_n)], \end{aligned}$$

where

$$u_k = f(X_k) - Pf(X_{k-1}) = f(X_k) - E(f(X_k) | X_{k-1})$$

are martingale differences with respect to  $\mathcal{F}_k = \sigma(X_0, X_1, \dots, X_k)$ , whereas

$$n^{-1/2} [Pf(X_0) - Pf(X_n)] \xrightarrow{P} 0$$

by Chebyshev's inequality. Thus, it is sufficient to prove that

$$Y_n := n^{-1/2} \sum_{k=1}^n u_k \xrightarrow{\mathcal{L}} N(0, \sigma^2)$$

and the convergence does not depend on the initial distribution. For this purpose, we shall show that the martingale differences {u<sub>k</sub>, k ≥ 1} satisfy the conditions (A<sub>1</sub>), (A<sub>2</sub>).

According to assumption (H<sub>2</sub>) we have

$$E_{\Pi}[E(u_1^2|\mathcal{F}_0)] = E_{\Pi}(u_1^2) = E_{\Pi}[f(X_1) - Pf(X_0)]^2 = E_{\Pi}f^2(X_1) - E_{\Pi}[Pf(X_0)]^2,$$

thus

$$E_{\Pi}(u_1^2) = \Pi f^2 - \Pi(Pf)^2 < \infty. \tag{9}$$

Therefore, by the ergodic Theorem 2.2, for any initial distribution with probability one

$$n^{-1} \sum_{k=1}^n E(u_k^2|\mathcal{F}_{k-1}) \longrightarrow E_{\Pi}u_1^2 = \sigma^2.$$

Thus the condition  $(A_1)$  of Theorem 2.3 is satisfied.

On the other hand, by (9) we have

$$E_{\Pi}(u_1^2 \mathbf{1}_{\{|u_1| \geq t\}}) \longrightarrow 0, \tag{10}$$

as  $t \uparrow \infty$ . Again by the ergodic Theorem 2.2, for any initial distribution, with probability one

$$n^{-1} \sum_{k=1}^n E(u_k^2 \mathbf{1}_{\{|u_k| \geq t\}}|\mathcal{F}_{k-1}) \longrightarrow E_{\Pi}(u_1^2 \mathbf{1}_{\{|u_1| \geq t\}}) \tag{11}$$

for each  $t > 0$ . By (11) and then (10) we have with probability one

$$\begin{aligned} 0 &\leq \overline{\lim}_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n E_{\Pi}(u_k^2 \mathbf{1}_{\{|u_k| \geq \varepsilon \sqrt{n}\}}) \\ &\leq \overline{\lim}_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n E_{\Pi}(u_k^2 \mathbf{1}_{\{|u_k| \geq t\}}) \\ &= E_{\Pi}(u_1^2 \mathbf{1}_{\{|u_1| \geq t\}}) \longrightarrow 0 \text{ as } t \uparrow \infty. \end{aligned}$$

Thus condition  $(A_2)$  is satisfied, hence by the central limit theorem for martingale differences  $\{u_k, k \geq 1\}$  (7) holds. ■

*Remark 2.* If the series

$$\sum_{n=0}^{\infty} P^n \varphi(x) = \sum_{n=0}^{\infty} \int_E \varphi(y) P^n(x, dy)$$

converges, then we always have

$$\varphi(x) = f(x) - Pf(x)$$

with

$$f(x) = \sum_{n=0}^{\infty} P^n \varphi(x).$$

In fact, it is obvious that

$$f(x) = \varphi(x) + \sum_{n=1}^{\infty} P^n \varphi(x) = \varphi(x) + P \sum_{n=0}^{\infty} P^n \varphi(x) = \varphi(x) + Pf(x).$$

Furthermore, in this case

$$\sigma^2 = \Pi \left[ \varphi^2 + 2 \sum_{n=0}^{\infty} \varphi P^n \varphi \right].$$

Remark 3. If  $\varphi = f - Pf$  holds, then

$$\Pi \varphi = \Pi f - \Pi Pf = 0. \tag{12}$$

So the condition (12) is necessary for  $\varphi = f - Pf$ . Furthermore, in addition if we have

$$\lim_{n \rightarrow \infty} P^n f(x) = \Pi f, \quad \forall x \in E$$

then  $f(x)$  is also given by

$$f(x) = \sum_{n=0}^{\infty} P^n \varphi(x) + \Pi f.$$

In fact, we have

$$\begin{aligned} \varphi(x) &= f(x) - Pf(x) \\ P\varphi(x) &= Pf(x) - P^2f(x) \\ &\dots \\ P^n \varphi(x) &= P^n f(x) - P^{n+1} f(x). \end{aligned}$$

Summing the above equalities we obtain

$$\sum_{k=0}^n P^k \varphi(x) = f(x) - P^{n+1} f(x) \longrightarrow f(x) - \Pi f.$$

Remark 4. Function  $f$  given by (6) is defined uniquely up to an additional constant if  $\lim_{n \rightarrow \infty} P^n g(x) = \Pi g$  for all  $g$   $\Pi$ -integrable.

In fact, suppose that  $f_1, f_2$  are the functions satisfying (6). Then  $g = f_1 - f_2$  is a solution of the equations:

$$g(x) = Pg(x), \quad g(x) = P(Pg(x)) = P^2g(x) = \dots = P^n g(x), \quad \forall x \in E$$

for all  $n = 1, 2, \dots$ . Thus there exists the limit

$$g(x) = \lim_{n \rightarrow \infty} P^n g(x) = \Pi g \text{ (a constant).}$$

It also follows from Remark 4 and from (8) that if  $f$  satisfies the equation (6) then  $\sigma^2$  is defined uniquely, i.e.,  $\sigma^2$  does not change if  $f$  is replaced by  $f + C$  with  $C$  being any constant, since



$$\Pi[\varphi^2 + 2\varphi P(f + C)] = \Pi[\varphi^2 + 2\varphi Pf] + 2C\Pi\varphi = \Pi[\varphi^2 + 2\varphi Pf].$$

*Remark 5.* If  $\Pi\varphi \neq 0$  we can replace  $\varphi$  by  $\varphi^* = \varphi - \Pi\varphi$ .

**Corollary 2.1.** *Assume that a Markov chain  $\{X_n, n \geq 0\}$  is irreducible, ergodic with the countable state space  $E = \{1, 2, \dots\}$  and with the ergodic distribution  $\Pi = (\pi_1, \pi_2, \dots)$  and the following condition is satisfied*  
*(H<sub>3</sub>) The mapping  $\varphi : E \rightarrow \mathbb{R}$  takes the form*

$$\varphi(x) = f(x) - Pf(x), \quad \forall x \in E$$

with  $f : E \rightarrow \mathbb{R}$  being measurable such that  $\Pi f^2 < \infty$ . Put

$$\sigma^2 = \Pi[f^2 - (Pf)^2] = \Pi[\varphi^2 + 2\varphi Pf].$$

Then

$$n^{-1/2} \sum_{k=1}^n \varphi(X_k) \xrightarrow{\mathcal{L}} N(0, \sigma^2) \text{ as } n \rightarrow \infty.$$

### 3. Central Limit for Integral Functional of Jump Markov Process

#### 3.1. Jump Markov Process

Let  $\{X(t), t \geq 0\}$  be a random process defined on some probability space  $(\Omega, \mathcal{F}, P)$  with measurable state space  $(E, \mathcal{B})$ .

**Definition 3.1.** *The process  $\{X(t), t \geq 0\}$  is called jump homogeneous Markov process with the state space  $(E, \mathcal{B})$  if it is a Markov process with transition probability*

$$P(t, x, A) = P(X(t+s) \in A | X(s) = x), \quad s, t \geq 0$$

satisfying the following condition

$$\lim_{t \rightarrow 0} P(t, x, \{x\}) = 1, \quad \forall x \in E. \quad (13)$$

We suppose also that  $\{X(t), t \geq 0\}$  is right continuous and the limit (13) is uniform in  $x \in E$ .

By Theorem 2.4 in [6] the sample functions of  $\{X(t), t \geq 0\}$  are step functions with probability one, and there exist two  $q$ -functions  $q(\cdot)$  and  $q(\cdot, \cdot)$  being Baire functions where  $q(x, \cdot)$  is finite measure on Borel subsets of  $E \setminus \{x\}$ ,  $q(x) = q(x, E \setminus \{x\})$  is bounded. Further

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{(1 - P(t, x, \{x\}))}{t} &= q(x), \\ \lim_{t \rightarrow 0} \frac{P(t, x, A)}{t} &= q(x, A) \end{aligned}$$

uniformly in  $A \subset E \setminus \{x\}$ .

If  $q(x) > 0 \forall x \in E$  then the process has no absorbing state. We assume also that  $q(x)$  is bounded from 0.

Since  $\{X(t), t \geq 0\}$  is right continuous and step process, the system starts out in some state  $Z_1$ , stays there a length of time  $\rho_1$ , then jumps immediately to a new state  $Z_2$ , stays a length of time  $\rho_2$ , etc. Therefore there exist random variables  $Z_1, Z_2, \dots$  and  $\rho_1, \rho_2, \dots$  such that

$$\begin{aligned} X(t) &= Z_1, \quad \text{if } 0 \leq t < \rho_1, \\ X(t) &= Z_n, \quad \text{if } \rho_1 + \dots + \rho_{n-1} \leq t < \rho_1 + \dots + \rho_n, \quad n \geq 2. \end{aligned}$$

$\rho_n$ 's are all finite because we have assumed that  $q(x) > 0 \forall x \in E$ .

Let  $\nu(t)$  be the random variable defined by

$$\nu(t) = \max\{k : \rho_1 + \dots + \rho_k < t\}$$

then  $\nu(t)$  is the number of jumps which occur up to time  $t$ .

It follows from the general theory of discontinuous Markov process (see [6], p.266) that  $\{Z_n, n \geq 1\}$  is a Markov chain with transition probability

$$P(x, A) = \frac{q(x, A)}{q(x)}, \tag{14}$$

furthermore

$$P(\rho_{n+1} > s | \rho_1, \dots, \rho_n, Z_1, \dots, Z_{n+1}) = e^{-q(Z_{n+1})s}, \quad s > 0 \tag{15}$$

$$P(Z_{n+1} \in A | \rho_1, \dots, \rho_n, Z_1, \dots, Z_n) = P(Z_n, A). \tag{16}$$

The function  $q(.,.)$  is called the transition intensity.

It follows from (15), (16) that  $\{(Z_n, \rho_n), n \geq 1\}$  is a Markov chain on the cartesian product  $E \times \mathbb{R}^+$ , where  $\mathbb{R}^+ = (0, \infty)$ . This chain is called the imbedded chain with the transition probability

$$\begin{aligned} Q(x, s, A \times B) &= P(Z_{n+1} \in A, \rho_{n+1} \in B | Z_n = x, \rho_n = s) \\ &= \int_A P(x, dy) \int_B q(y) e^{-q(y)u} du, \end{aligned}$$

$A \times B \in \mathcal{B} \times \mathcal{B}(\mathbb{R}^+)$ , where  $\mathcal{B}(\mathbb{R}^+)$  denotes the Borel  $\sigma$ - algebra on  $\mathbb{R}^+$ . This transition probability does not depend on  $s$  and we rewrite it by  $Q(x, A \times B)$  or formally by

$$Q(x, dy \times du) = P(x, dy)q(y) \exp(-q(y)u)du.$$

**Definition 3.2.** The probability measure  $\Pi^*$  on  $(E \times \mathbb{R}^+, \mathcal{B} \times \mathcal{B}(\mathbb{R}^+))$  is called the stationary distribution of the imbedded chain  $\{(Z_n, \rho_n), n \geq 1\}$  if

$$\Pi^*(A \times B) = \int_{E \times \mathbb{R}^+} \Pi^*(dx \times ds)Q(x, A \times B), \quad A \times B \in \mathcal{B} \times \mathcal{B}(\mathbb{R}^+). \tag{17}$$

Letting  $B = \mathbb{R}^+$ , then  $\Pi^*$  is the stationary distribution of the imbedded chain if and only if

$$\Pi(\cdot) = \Pi^*(\cdot \times \mathbb{R}^+) \quad (18)$$

is the one of  $\{Z_n, n \geq 1\}$  with the transition probability  $P(x, A) = Q(x, A \times \mathbb{R}^+)$  and

$$\Pi^*(A \times B) = \int_E \Pi(dx) Q(x, A \times B).$$

Since  $\Pi P(\cdot) = \Pi(\cdot)$ , we have

$$\begin{aligned} \Pi^*(A \times B) &= \int_E \Pi(dx) \int_A P(x, dy) \int_B q(y) \exp(-q(y)u) du \\ &= \int_A \left( \int_E \Pi(dx) P(x, dy) \right) \int_B q(y) \exp(-q(y)u) du \end{aligned}$$

or

$$\Pi^*(A \times B) = \int_A \Pi(dy) \int_B q(y) \exp(-q(y)u) du \quad (19)$$

or in differential form

$$\Pi^*(dy \times du) = \Pi(dy) q(y) \exp(-q(y)u) du. \quad (20)$$

Thus we have the following proposition:

**Proposition 3.1.** *If the Markov chain  $\{Z_n, n \geq 1\}$  with the transition probability  $P(x, A)$  has the stationary distribution  $\Pi$  then the imbedded chain possesses also the stationary distribution  $\Pi^*$  defined by (19) or (20).*

**Proposition 3.2.** *If  $P(x, \cdot) \ll \Pi(\cdot) \forall x \in E$ , where  $\Pi$  is the stationary distribution of  $\{Z_n, n \geq 1\}$  then the transition probability  $Q(x, \cdot)$  of the imbedded chain is also absolutely continuous with respect to the stationary distribution  $\Pi^*$ , i.e.*

$$Q(x, \cdot) \ll \Pi^*(\cdot), \forall x \in E.$$

(see [3], p.66).

Here and after we shall denote by  $\Pi, \Pi^*$  the stationary distributions of Markov chain  $\{Z_n, n \geq 1\}$  and the imbedded chain  $\{(Z_n, \rho_n), n \geq 1\}$ , respectively.

### 3.2. Functional Central Limit Theorem

We have the following ergodic theorem for the imbedded chain

**Theorem 3.1.** (Ergodic theorem for the imbedded process) *If Markov chain  $\{Z_n, n \geq 1\}$  with the transition probability  $P(x, \cdot)$  having the stationary distribution  $\Pi$  such that*

$$P(x, \cdot) \ll \Pi(\cdot) \quad \forall x \in E,$$

and if  $\varphi(Z_1, \rho_1; Z_2, \rho_2)$  is the random variable possessing the finite expectation  $\mu$  w.r.t. the probability measure  $P_{\Pi^*}$ , then for any initial distribution

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n \varphi(Z_k, \rho_k; Z_{k+1}, \rho_{k+1}) = \mu; \text{ a.s.} \tag{21}$$

In particular, if  $\Pi q^{-1} < \infty$  then

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n \rho_k = \int_E \Pi(dy)(q(y))^{-1} = \Pi q^{-1} \text{ a.s.} \tag{22}$$

Furthermore

$$\lim_{t \rightarrow \infty} \frac{\nu(t)}{t} = (\Pi q^{-1})^{-1} =: \alpha > 0 \text{ a.s.} \tag{23}$$

and (21), (22) remain valid if in the limits  $n$  is replaced by  $\nu(t)$ , then limits are taken as  $t \rightarrow \infty$ .

*Proof.* (21) follows from the ergodic theorem for Markov chain  $\{(Z_n, \rho_n), n \geq 1\}$ , and (23) follows from (22) by the same argument as in the renewal theory. ■

Applying Theorem 2.4 for the imbedded chain  $\{(Z_n, \rho_n), n \geq 1\}$  we obtain the following theorem.

**Theorem 3.2.** (Central limit theorem for the imbedded chain) *Assume that the following conditions  $(C_1), (C_2)$  are satisfied:*

$(C_1)$  *The jump Markov process  $\{X(t), t \geq 0\}$  has the imbedded chain  $\{(Z_n, \rho_n), n \geq 1\}$  such that the Markov chain  $\{Z_n, n \geq 1\}$  has the transition probability  $P(x, \cdot)$  with the stationary distribution  $\Pi$  satisfying the following condition*

$$P(x, \cdot) \ll \Pi(\cdot) \quad \forall x \in E.$$

$(C_2)$  *The function  $\psi : E \times \mathbb{R}^+ \rightarrow \mathbb{R}$  takes the form*

$$\psi(x, s) = f(x, s) - Qf(x, s),$$

where  $f : E \times \mathbb{R}^+ \rightarrow \mathbb{R}$  is  $\mathcal{B} \times \mathcal{B}(\mathbb{R}^+)$ -measurable and

$$Qf(x) = Qf(x, s) = \int_E P(x, dy) \int_{\mathbb{R}^+} f(y, u)q(y) \exp(-q(y)u)du.$$

Furthermore, the function  $f$  has the following property

$$\Pi^* f^2 = \int_E \Pi(dy) \int_{\mathbb{R}^+} |f(y, u)|^2 q(y) \exp(-q(y)u)du < \infty. \tag{24}$$

Then we have

$$n^{-1/2} \sum_{k=1}^n \psi(Z_k, \rho_k) \xrightarrow{\mathcal{L}} N(0, \sigma^2) \tag{25}$$

for any initial distribution, where

$$\sigma^2 = \Pi^*(f^2 - (Qf)^2) = \Pi^*(\psi^2 + 2\psi Qf). \tag{26}$$

The goal of this section is to investigate the limit law of the integral functional  $T^{-1/2} \int_0^T \varphi(X(t))dt$  as  $T \rightarrow \infty$ .

Let us at first notice that

$$\int_0^T \varphi(X(t))dt = \sum_{k=1}^{\nu(T)} \varphi(Z_k)\rho_k + \varphi(Z_{\nu(T)+1})(T - \tau_{\nu(T)}), \tag{27}$$

where

$$\tau_1 = \rho_1, \tau_2 = \rho_1 + \rho_2, \dots, \tau_n = \rho_1 + \rho_2 + \dots + \rho_n, \dots$$

are the jump times of the process  $\{X(t), t \geq 0\}$ .

In what follows we suppose always that the condition  $(C_1)$  is satisfied.

We need the following lemmas.

**Lemma 3.1.** *If  $\Pi\varphi^2q^{-2} < \infty$  then*

$$\frac{1}{\sqrt{T}}\varphi(Z_{\nu(T)+1})(T - \tau_{\nu(T)}) \xrightarrow{P} 0 \tag{28}$$

for any initial distribution.

*Proof.* Noticing that for  $\psi(x, s) = \varphi(x)s$  we have

$$\begin{aligned} \Pi^*\psi^2 &= \int_E \Pi(dy)\varphi^2(y) \int_{\mathbb{R}^+} u^2q(y) \exp(-q(y)u)du \\ &= 2 \int_E \varphi^2(y)q^{-2}(y)\Pi(dy) = 2\Pi\varphi^2q^{-2} < \infty \end{aligned}$$

and  $\nu(T) \rightarrow \infty$  a.s. as  $T \rightarrow \infty$  by (23). By those and by the ergodic Theorem 3.1

$$(\nu(T) + 1)^{-1} \sum_{k=1}^{\nu(T)+1} |\varphi(Z_k)\rho_k|^2 \longrightarrow \Pi^*\psi^2 \text{ a.s.}$$

as  $T \rightarrow \infty$ . Hence with probability one

$$(\nu(T) + 1)^{-1}|\varphi(Z_{\nu(T)+1})\rho_{\nu(T)+1}|^2 \longrightarrow 0$$

and (28) follows from

$$\frac{1}{\sqrt{T}}|\varphi(Z_{\nu(T)+1})(T - \tau_{\nu(T)})| \leq \left(\frac{\nu(T) + 1}{T}\right)^{1/2}(\nu(T)+1)^{-1/2}|\varphi(Z_{\nu(T)+1})\rho_{\nu(T)+1}| \rightarrow 0$$

a.s. ■

**Lemma 3.2.** *Suppose that  $\{u_k, \mathcal{F}_k, k \geq 1\}$  defined on  $(\Omega, \mathcal{F}, P)$  are the square integrable martingale differences such that*

$$\sup_{n, m \geq 1} (n^{-1} \sum_{k=m}^{m+n} E u_k^2) = C < \infty \tag{29}$$

and that  $\{\nu(t), t \geq 0\}$  is a random process valued in  $\{1, 2, \dots\}$  such that  $\{\nu(t) = k\} \in \mathcal{F}_k \forall t \geq 0$  and

$$\lim_{t \rightarrow \infty} \frac{\nu(t)}{t} = \alpha > 0 \text{ a.s.} \tag{30}$$

Then

$$T^{-1/2} \left| \sum_{k=1}^{\nu(T)} u_k - \sum_{k=1}^{[\alpha T]} u_k \right| \xrightarrow{P} 0 \text{ as } T \rightarrow \infty. \tag{31}$$

*Proof.* It follows from condition (30) that: for all  $\varepsilon > 0$ , and  $T$  sufficiently large we have

$$P\left(\left|\frac{\nu(T)}{T} - \alpha\right| > \varepsilon^3\right) < \varepsilon$$

or

$$P((\alpha - \varepsilon^3)T < \nu(T) < (\alpha + \varepsilon^3)T) \geq 1 - \varepsilon. \tag{32}$$

Putting

$$A_\varepsilon = \{(\alpha - \varepsilon^3)T < \nu(T) < (\alpha + \varepsilon^3)T\},$$

we have

$$\begin{aligned} P\left(T^{-\frac{1}{2}} \left| \sum_{k=1}^{\nu(T)} u_k - \sum_{k=1}^{[\alpha T]} u_k \right| > \varepsilon\right) &\leq P(A_\varepsilon^c) + P\left(\left\{T^{-\frac{1}{2}} \left| \sum_{k=1}^{\nu(T)} u_k - \sum_{k=1}^{[\alpha T]} u_k \right| > \varepsilon\right\} \cap A_\varepsilon\right) \\ &\leq \varepsilon + P\left(T^{-\frac{1}{2}} \max_{|l - [\alpha T]| < \varepsilon^3 T} \left| \sum_{k=1}^l u_k - \sum_{k=1}^{[\alpha T]} u_k \right| > \varepsilon\right) \\ &\leq \varepsilon + P\left(\max_{a \leq l \leq b} \left| \sum_{k=a}^l u_k \right| > \frac{\varepsilon T^{\frac{1}{2}}}{2}\right) \end{aligned} \tag{33}$$

where  $a = [\alpha T] - [\varepsilon^3 T]$ ,  $b = [\alpha T] + [\varepsilon^3 T]$  with  $[r]$  denoting the integer part of the number  $r$ .

By Kolmogorov's inequality for a martingale

$$P\left(\max_{1 \leq n \leq N} \left| \sum_{k=1}^n u_k \right| > \lambda\right) \leq \frac{1}{\lambda^2} E \left[ \sum_{k=1}^N u_k \right]^2 = \frac{1}{\lambda^2} \sum_{k=1}^N E u_k^2,$$

we have

$$P\left(\max_{a \leq l \leq b} \left| \sum_{k=a}^l u_k \right| > \frac{\varepsilon T^{\frac{1}{2}}}{2}\right) \leq 8\varepsilon \frac{1}{2\varepsilon^3 T} E \left[ \sum_{k=a}^{2[\varepsilon^3] + a} u_k^2 \right] \leq 8\varepsilon C. \tag{34}$$

It follows from (33), (34) that (31) holds. ■

**Corollary 3.1.** *Assume that the martingale differences  $\{u_k, k \geq 1\}$  take the form*

$$u_k = f(X_k) - E(f(X_k)|X_{k-1}), \quad k = 1, 2, \dots$$

where  $\{X_k, k \geq 0\}$  is a Markov chain with the stationary distribution  $\Pi$  such that  $\Pi f^2 < \infty$ . Then (31) holds for any initial distribution.

*Proof.* It is obvious that

$$E_{\Pi} u_k^2 \leq E_{\Pi} f^2(X_{k-1}) = \Pi f^2 < \infty,$$

therefore

$$E_{\Pi}(n^{-1} \sum_{k=m}^{m+n} u_k^2) \leq \Pi f^2 = C, \quad \forall m, n.$$

Denoting the quantity in the left-hand side of (31) by  $\eta_T$ , by Lemma 3.2 we obtain

$$\lim_{T \rightarrow \infty} P_{\Pi}(|\eta_T| \geq \varepsilon) = 0 \quad \forall \varepsilon > 0$$

or

$$\lim_{T \rightarrow \infty} \int_E P_x(|\eta_T| \geq \varepsilon) \Pi(dx) = 0 \quad \forall \varepsilon > 0.$$

It follows that there exists a subset  $\Lambda \subset E$  such that  $\Pi(\Lambda) = 0$  and

$$\lim_{T \rightarrow \infty} P_x(|\eta_T| \geq \varepsilon) = 0 \quad \forall x \in E \setminus \Lambda.$$

Since  $P(x, \cdot) \ll \Pi(\cdot) \quad \forall x, P(x, E \setminus \Lambda) = 1 \quad \forall x \in E$ .

On the other hand, letting  $A_T = \{|\eta_T| \geq \varepsilon\}$ , we observe that  $A_T \in \cup_{n \geq n_0} \mathcal{F}_n$  with  $n_0 > 1$ , where  $\mathcal{F}_n = \sigma(X_k, k \geq n)$ . Then by Markov property:

$$\begin{aligned} P_x(A_T) &= E(1_{A_T} | X_0 = x) = E[E(1_{A_T} | X_1) | X_0 = x] \\ &= \int_E E(1_{A_T} | X_1 = y) P(x, dy) = \int_{E \setminus \Lambda} E_y(1_{A_T}) P(x, dy). \end{aligned}$$

Therefore

$$\begin{aligned} 0 \leq \limsup_{T \rightarrow \infty} P_x(A_T) &= \limsup_{T \rightarrow \infty} \int_{E \setminus \Lambda} P_y(A_T) P(x, dy) \\ &= \int_{E \setminus \Lambda} \lim_{T \rightarrow \infty} P_y(A_T) P(x, dy) = 0. \end{aligned}$$

So

$$\lim_{T \rightarrow \infty} P_x(A_T) = 0 \quad \forall x$$

and hence

$$\lim_{T \rightarrow \infty} P_{\nu}(|\eta_T| \geq \varepsilon) = \lim_{T \rightarrow \infty} \int_E P_x(|\eta_T| \geq \varepsilon) \nu(dx) = 0.$$

This implies (31). ■

**Lemma 3.3.** Assume that the following equation has a solution  $g(x)$

$$(I - P)g(x) = P\varphi q^{-1}(x). \quad (35)$$

Then, putting

$$f(x, s) = \varphi(x)s + g(x), \quad (36)$$

we have the representation

$$\varphi(x)s = f(x, s) - Qf(x), \quad (37)$$

where  $Qf(x) = g(x)$ .

*Proof.* At first let us notice that for  $\psi : E \times \mathbb{R}^+ \rightarrow \mathbb{R}$  given by  $\psi(x, s) = \varphi(x)s$  we have

$$\begin{aligned} Qg(x) &= \int_E g(y)P(x, dy) \int_{\mathbb{R}^+} q(y) \exp(-q(y)u) du \\ &= \int_E g(y)P(x, dy) = Pg(x), \\ Q\psi(x) &= \int_E \varphi(y)P(x, dy) \int_{\mathbb{R}^+} uq(y) \exp(-q(y)u) du \\ &= \int_E \varphi(y)q^{-1}(y)P(x, dy) = P\varphi q^{-1}(x). \end{aligned}$$

In order to prove (37) we shall prove that if  $g(x)$  is a solution of (35) then  $g(x) = Qf(x)$ . In fact, by (36)

$$Qf(x) = Q\psi(x) + Qg(x) = P\varphi q^{-1}(x) + Pg(x) = g(x). \quad \blacksquare$$

*Remark 6.* A necessary condition for the existence of a solution of (35) is

$$\Pi\varphi q^{-1} = 0. \quad (38)$$

In fact, applying operator  $\Pi$  on both sides of (35) we have  $\Pi g - \Pi P g = 0 = \Pi\varphi q^{-1}$ .

Let us notice that the condition (38) is satisfied if the function  $\varphi$  is represented in the form

$$\varphi(x) = \varphi^*(x) - \alpha \Pi\varphi^* q^{-1}$$

where  $\varphi^* : E \rightarrow \mathbb{R}$ ,  $\alpha$  is given by (23).

**Lemma 3.4.** Assume that the following equation has a solution  $g$

$$(I - P)g(x) = P\varphi q^{-1}(x)$$

and that  $\Pi\varphi^2 q^{-2} < \infty$ ,  $\Pi g^2 < \infty$ . Furthermore, if the condition  $(C_1)$  of Theorem 3.2 are satisfied then



$$T^{-1/2} \sum_{k=1}^{\nu(T)} \varphi(Z_k) \rho_k \xrightarrow{\mathcal{L}} N(0, \alpha \delta^2) \tag{39}$$

for any initial distribution, where  $\alpha$  is given by (23) and

$$\delta^2 = 2\Pi(\varphi^2 q^{-2} + \varphi q^{-1} g).$$

*Proof.* By Lemma 3.3, we have the representation

$$\begin{aligned} \psi(Z_k, \rho_k) &= \varphi(Z_k) \rho_k = f(Z_k, \rho_k) - Qf(Z_k) \\ &= f(Z_k, \rho_k) - Qf(Z_{k-1}) + Qf(Z_{k-1}) - Qf(Z_k) \\ &= u_k + g(Z_{k-1}) - g(Z_k) \end{aligned}$$

where  $\{u_k = f(Z_k, \rho_k) - Qf(Z_{k-1}), k \geq 1\}$  are martingale differences. Therefore

$$\begin{aligned} T^{-1/2} \sum_{k=1}^{\nu(T)} \varphi(Z_k) \rho_k &= T^{-1/2} \sum_{k=1}^{\nu(T)} u_k + T^{-1/2} \sum_{k=1}^{\nu(T)} (g(Z_{k-1}) - g(Z_k)) \\ &= T^{-1/2} \sum_{k=1}^{\nu(T)} u_k + T^{-1/2} (g(Z_0) - g(Z_{\nu(T)})). \end{aligned} \tag{40}$$

Since  $\Pi g^2 < \infty$ , by the same argument as in the proof of Lemma 3.1, we can show that

$$T^{-1/2} (g(Z_0) - g(Z_{\nu(T)})) \xrightarrow{P} 0 \tag{41}$$

for any initial distribution.

Furthermore, we have by (36)

$$\Pi^* f^2 \leq 2\Pi^*(\psi^2 + g^2) = 2(\Pi\varphi^2 q^{-2} + \Pi g^2) < \infty,$$

hence by Corollary 3.1, (31) holds for any initial distribution.

Applying Theorem 3.2 for the imbedded chain  $\{(Z_k, \rho_k), k \geq 1\}$  we obtain

$$T^{-1/2} \sum_{k=1}^{[\alpha T]} u_k \xrightarrow{\mathcal{L}} N(0, \alpha \delta^2) \tag{42}$$

with

$$\begin{aligned} \delta^2 &= \Pi^*(f^2 - (Qf)^2) = \Pi^*(f^2 - g^2) \\ &= \Pi^*(\psi^2 + 2\psi g) = 2\Pi(\varphi^2 q^{-2} + \varphi g q^{-1}). \end{aligned}$$

Finally, it follows from (40), (31), (41), (34), (42) that (39) holds for any initial distribution. ■

Now we state and prove the main theorem as follows

**Theorem 3.3.** Assume that the condition  $(C_1)$  of Theorem 3.2 and the following condition  $(C_3)$  are satisfied

$(C_3)$  (i)  $\Pi\varphi^2q^{-2} < \infty$  and, (ii) The following equation has a solution  $g$

$$(I - P)g(x) = P\varphi q^{-1}(x)$$

with  $\Pi g^2 < \infty$ .

Then

$$T^{-1/2} \int_0^T \varphi(X(t))dt \xrightarrow{\mathcal{L}} N(0, \alpha\delta^2)$$

for any initial distribution, where

$$\delta^2 = 2\Pi(\varphi^2q^{-2} + \varphi gq^{-1}).$$

*Proof.* The conclusion of Theorem 3.3 follows from Theorem 3.2 and Lemmas 3.1, 3.4. ■

#### 4. Examples

*Example 1.* Assume that the jump Markov process  $\{X(t), t \geq 0\}$  with the state space  $E = \{1, 2, 3\}$  has the transition intensity matrix

$$Q = \begin{bmatrix} -1 & 0.5 & 0.5 \\ 0.4 & -1 & 0.6 \\ 0.8 & 0.2 & -1 \end{bmatrix}.$$

Then the Markov chain  $\{Z_k, k \geq 1\}$  has the transition probability matrix

$$P = \begin{bmatrix} 0 & 0.5 & 0.5 \\ 0.4 & 0 & 0.6 \\ 0.8 & 0.2 & 0 \end{bmatrix}.$$

It is easy to see that  $\{Z_k, k \geq 1\}$  possesses the ergodic distribution as follows

$$\Pi = [0.38596 \quad 0.26316 \quad 0.35088],$$

whereas the sequence  $\{\rho_k, k \geq 1\}$  is the sequence of independent, exponentially indentially distributed random variables with the parameter  $q = 1$  (i.e.,  $q(x) = 1$  for all  $x \in E$ ) and hence  $\alpha = 1$ .

Let us consider  $\varphi^* = [1, 2, 4]^T$ , i.e.  $\varphi^*(1) = 1, \varphi^*(2) = 2, \varphi^*(3) = 4$ . Then

$$\Pi\varphi^* = 2.3158, \quad \varphi = \varphi^* - \Pi\varphi^* = [-1.3158 \quad -0.3158 \quad 1.6842]^T.$$

We shall prove that as  $T \rightarrow \infty$

$$\frac{1}{\sqrt{T}} \int_0^T (\varphi^*(X(t)) - 2.3158)dt \xrightarrow{\mathcal{L}} N(0, \sigma^2). \quad (43)$$

For this purpose, we try to find a function  $g = [g_1, g_2, g_3]^T$  satisfying the following equation

$$(I - P)g = P\varphi q^{-1} = P\varphi$$

or in detail

$$\begin{bmatrix} 1 & -0.5 & -0.5 \\ -0.4 & 1 & -0.6 \\ -0.8 & -0.2 & 1 \end{bmatrix} \begin{bmatrix} g_1 \\ g_2 \\ g_3 \end{bmatrix} = \begin{bmatrix} 0.6842 \\ 0.4842 \\ -1.1158 \end{bmatrix}.$$

The above algebraic equation has the following solution

$$g = [1.15788 \quad 0.94735 \quad 0].$$

Since  $E$  has a finite number of elements,  $\Pi g^2$  and  $\Pi\varphi^2$  are finite. By Theorem 3.3 we have (43) with

$$\sigma^2 = \delta^2 = 2\Pi(\varphi^2 + \varphi g) = 2.046.$$

*Example 2.* Let us consider the integral functional of the jump Markov process with the state space  $E = \{1, 2, \dots\}$  defined by:

$$T_i = \int_0^T \mathbf{1}_{\{X(t)=i\}}, \quad i \in E.$$

This integral is the total time length during which the process visits the state  $i$ . Assume that this process satisfies the condition  $(C_1)$ .

For each state  $i$ , put  $\varphi^*(x) = \mathbf{1}_{\{x=i\}}$  then  $\alpha\Pi\varphi^*q^{-1} = \alpha\pi_i q_i^{-1}$ . Let us consider  $\varphi(x) = \varphi^*(x) - \alpha\pi_i q_i^{-1}$ .

Suppose that the equation

$$(I - P)g(x) = P\varphi q^{-1}(x) \tag{44}$$

has a solution  $g$  such that  $\Pi g^2 < \infty$ . Then by Theorem 3.3

$$\frac{1}{\sqrt{T}} \int_0^T \varphi(X(t))dt = \frac{1}{\sqrt{T}} \int_0^T (\mathbf{1}_{\{X(t)=i\}} - \alpha\pi_i q_i^{-1})dt \xrightarrow{\mathcal{L}} N(0, \alpha\delta^2)$$

where  $\delta^2 = 2\Pi(\varphi^2 q^{-2} + \varphi q^{-1}g)$ .

In particular, for the case where

$$E = \{1, 2\}, \quad Q = \begin{bmatrix} -q_1 & q_1 \\ q_2 & -q_2 \end{bmatrix}, \quad P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad q_1, q_2 > 0,$$

we have the stationary distribution  $\Pi = (1/2, 1/2)$  and

$$\alpha = (\Pi q^{-1})^{-1} = \left(\frac{1}{2}\left(\frac{1}{q_1} + \frac{1}{q_2}\right)\right)^{-1} = \frac{2q_1 q_2}{q_1 + q_2}.$$

Put  $\varphi^*(x) = \mathbf{1}_{\{x=1\}}$ , then

$$\alpha\Pi\varphi^*q^{-1} = \alpha\pi_1 q_1^{-1} = \frac{q_2}{q_1 + q_2}, \quad \varphi(x) = \mathbf{1}_{\{x=1\}} - \frac{q_2}{q_1 + q_2},$$

and

$$\frac{1}{\sqrt{T}} \int_0^T \left(\mathbf{1}_{\{X(t)=1\}} - \frac{q_2}{q_1 + q_2}\right)dt \xrightarrow{\mathcal{L}} N(0, \alpha\delta^2) \tag{45}$$

for any initial distribution. In order to find  $\delta^2$  we have to solve the equation (44) for  $i = 1$ , i.e.

$$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}; \begin{bmatrix} g_1 \\ g_2 \end{bmatrix} = \begin{bmatrix} P\varphi q^{-1}(1) \\ P\varphi q^{-1}(2) \end{bmatrix} \quad (46)$$

with notice that

$$\begin{bmatrix} P\varphi q^{-1}(1) \\ P\varphi q^{-1}(2) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} (1 - q_2/(q_1 + q_2))q_1^{-1} \\ -(q_2/(q_1 + q_2))q_2^{-1} \end{bmatrix} = \begin{bmatrix} -1/(q_1 + q_2) \\ 1/(q_1 + q_2) \end{bmatrix}.$$

(46) has a solution  $g_1 = -1/(q_1 + q_2)$ ,  $g_2 = 0$ . Hence, by Theorem 3.3, we obtain (45) with

$$\delta^2 = 2\Pi(\varphi^2 q^{-2} + \varphi q^{-1}g) = \frac{1}{(q_1 + q_2)^2}.$$

We obtain from (45)

$$\sqrt{T} \left( \frac{T_1}{T} - \frac{q_2}{(q_1 + q_2)} \right) \xrightarrow{\mathcal{L}} N \left( 0, \frac{2q_1 q_2}{(q_1 + q_2)^3} \right).$$

*Acknowledgement.* The authors would like to thank Prof. Dr. Nguyen Huu Du and Prof. Dr. Tran Hung Thao for useful discussions.

## References

1. A. de Acosta, Moderate deviations for vector valued functional of a Markov chain: Lower bounds, *Ann. Probab.* **25** (1997) 259–294.
2. Athreya and P. Ney, A new approach to the limit theory of recurrent Markov chain, *Trans. Amer. Math. Society* **245** (1978) 493–501.
3. P. Billingsley, *Statistical Inference for Markov Processes*, The University of Chicago Press, 1958.
4. X. Chen, Limit theorem for functionals of Ergodic Markov Chains with general state space, *Memoir of the American Mathematical Society* **139** (1999) 1–200.
5. K. L. Chung, *Markov Chains with Stationary Transition Probabilities*, Springer-Verlag, Berlin, 2nd Edition, 1967.
6. J. L. Doob, *Stochastic Process*, John Wiley & Sons, New York, 1953.
7. P. Hall and C. C. Heyde, *Martingale Limit Theory and Its Application*, Academic Press, 1980.
8. S. Niem and E. Nummelin, *Central Limit Theorems for Markov Random Walks*, Commentations Physico-Mathematica, 54, Societas Scientiarum Fennica, Helsinki, 1982.
9. S. P. Meyn and R. L. Tweedie, *Markov Chains and Stochastic Stability*, Springer-Verlag, London, 1993.
10. E. Nummelin, *General Irreducible Markov Chains and Non-negative Operators*, Cambridge University Press, 1984.
11. M. S. Ross, *Introduction to Probability Models*, 7th Edition, Harcourt Academic Press, 2000.

# Vietnam Journal of MATHEMATICS

Volume 33, Number 4 December 2005

## CONTENTS FOR VOL. 33, NUMBER 4/2005 VIETNAM JOURNAL OF MATHEMATICS

A Characterization of Morrey Type Besov and Triebel-Lizorkin Spaces <b>Jingshi Xu</b>	369
On a System of Semilinear Elliptic Equations on an Unbounded Domain <b>Hoang Quoc Toan</b>	381
Mountain Pass Theorem and Nonuniformly Elliptic Equations <b>Nguyen Thanh Vu</b>	391
The Model of Stochastic Control and Applications <b>Nguyen Hong Hai and Dang Thanh Hai</b>	409
On the Generalized Convolution with a Weight-Function for Fourier, Fourier Cosine and Sine Transforms <b>Nguyen Xuan Thao and Nguyen Minh Khoa</b>	421
A Stochastic EOQ Policy of Cold-Drink-for a Retailer <b>S. S. Sana and K. Chaudhuri</b>	437
Central Limit Theorem for Functional of Jump Markov Processes <b>Nguyen Van Huu, Vuong Quan Hoang, and Tran Minh Ngoc</b>	443
On Efficient Sets in $\mathbb{R}^2$ <b>Hoang Xuan Phu</b>	463
Stress Field in an Elastic Strip in Frictionless Contact with a Rigid Stamp <b>Dang Dinh Ang</b>	469
On the Parametric Affine Variational Inequality Approach to Linear Fractional Vector Optimization Problems <b>T. N. Hoa, T. D. Phuong, and N. D. Yen</b>	477
Contents for Volume 33	491